

"OPTIMIZATION OF ORTHODONTIC TREATMENT METHODS FOR PATIENTS WITH DISTAL OCCLUSION"

for the degree of Master in Dentistry

Candidate: **Rakhimjonova Kh.U.**

Scientific Advisor: PhD, **Akhrorova M.Sh.**

Abstract:

Distal occlusion, a common malocclusion characterized by a posterior relationship of the mandibular arch to the maxillary arch, presents significant challenges in orthodontic treatment. This study explores the optimization of various orthodontic methods aimed at correcting distal occlusion, focusing on effectiveness, efficiency, and patient outcomes. We reviewed and analyzed current treatment modalities, including fixed appliances, functional appliances, and surgical options, considering factors such as age, skeletal discrepancies, and patient compliance. The investigation identifies specific treatment strategies and techniques that demonstrate improved clinical results and reduced treatment duration. Furthermore, we address the long-term stability of outcomes and potential side effects. The results emphasize the need for personalized treatment plans, integrating a combination of techniques to achieve optimal and sustainable correction of distal occlusion. This research aims to contribute to evidence-based orthodontic practice, ultimately enhancing patient care and satisfaction.

Keywords: Distal Occlusion, Malocclusion, Orthodontic Treatment, Fixed Appliances, Functional Appliances, Orthognathic Surgery, Treatment Optimization, Cephalometric Analysis, Patient Outcomes, Stability, Personalized Treatment

Introduction

Distal occlusion, also known as Class II malocclusion, is a prevalent orthodontic condition characterized by a returned mandible relative to the maxilla. This malocclusion often presents with a variety of clinical features, including an increased overjet, deep bite, and a convex facial profile. Distal occlusion can have significant impacts on both the functional efficiency of the masticatory system and the aesthetic appearance of the individual, potentially affecting their self-esteem and overall quality of life. Addressing this complex malocclusion requires a thorough

understanding of the underlying skeletal and dental discrepancies, and the judicious application of appropriate orthodontic treatment modalities.

Traditional orthodontic approaches to managing distal occlusion have included growth modification techniques during adolescence, such as headgear and functional appliances, as well as camouflage treatment in adults with premolar extractions and fixed appliances. However, these traditional approaches have limitations. Growth modification is only effective during the active growth phase and may not always be predictable. Camouflage treatment, while addressing the dental aspects of the malocclusion, may not fully correct the underlying skeletal discrepancies, potentially compromising long-term stability and facial aesthetics. Moreover, conventional fixed appliances can be associated with increased treatment time and patient discomfort.

The advent of new technologies and biomechanical principles in orthodontics has led to a greater focus on optimizing treatment outcomes for patients with distal occlusion. These advancements include skeletal anchorage devices (TADs), self-ligating brackets, and clear aligners, which offer alternatives to traditional techniques. TADs can provide stable anchorage for skeletal and dental movement, potentially reducing reliance on patient compliance with extraoral appliances. Self-ligating brackets, with their reduced friction, have been associated with improved treatment efficiency, and clear aligners offer a more aesthetically pleasing alternative to conventional fixed appliances.

The optimal application of these newer treatment modalities in the management of distal occlusion requires a careful consideration of individual patient characteristics, including age, growth status, the severity of the malocclusion, and patient preferences. This study aims to investigate and evaluate the effectiveness of different contemporary orthodontic treatment methods for patients with distal occlusion, aiming to identify optimized treatment protocols that can improve treatment outcomes, reduce treatment time, and enhance overall patient satisfaction. The study will focus on comparing traditional approaches with newer techniques, exploring the potential of these innovations to address the skeletal, dental, and soft tissue challenges associated with distal occlusion, striving to contribute to evidence-based practice in the management of distal occlusion.

Materials and Method

1. Traditional Treatment Group (Control): Patients in this group will be treated using traditional orthodontic methods, which may include:

- Growth modification appliances (e.g., headgear, functional appliances) during adolescence, if applicable
- Premolar extraction and fixed appliances.

2. TADs-Based Treatment Group: Patients in this group will be treated using fixed appliances combined with Temporary Anchorage Devices (TADs), aiming to achieve skeletal correction and controlled tooth movement.

3. Clear Aligner Treatment Group: Patients in this group will be treated using clear aligners for dental correction, where appropriate based on the severity of the malocclusion. This may or may not include adjunct therapy to aid in skeletal corrections if needed.

4. Combined Approach Group: This group may get treatment with a combination of the above techniques based on the treatment needs of individual patients.

Data Collection:

Data will be collected at the following time points:

- Baseline (T1): Pre-treatment data will be collected including:
 - Comprehensive orthodontic records including panoramic and cephalometric radiographs.
 - Intraoral and extraoral photographs
 - Dental casts.
 - Patient-reported outcome measures (PROMs), such as questionnaires on oral health-related quality of life and patient satisfaction.
- Mid-Treatment (T2): Progress records will be taken at predetermined intervals, based on treatment progress, including:
 - Intraoral photographs.
 - Cephalometric radiographs will be taken where clinically relevant and per individual treatment needs.
- Post-Treatment (T3): Post-treatment data will be collected including:

- Comprehensive orthodontic records, similar to baseline (T1).
- PROMs.

Outcome Measures:

The following outcome measures will be assessed:

- **Skeletal Changes:** Cephalometric measurements will be used to assess changes in skeletal relationships.
- **Dental Changes:** Linear and angular measurements will be made to assess changes in overjet, overbite, molar relationship, and tooth alignment.
- **Soft Tissue Changes:** Changes in facial profile and lip position will be evaluated using photographs.
- **Treatment Duration:** The duration of active orthodontic treatment will be recorded.
- **Patient-Reported Outcomes:** PROMs will be used to evaluate patient satisfaction, oral health-related quality of life, and treatment-related discomfort.
- **Treatment Stability:** Patients will be followed over a year after active treatment to determine treatment stability, and records will be taken at these follow-up appointments to monitor relapse if any.

Data Analysis:

Data will be analyzed using appropriate statistical methods, including ANOVA and t-tests, to compare the changes in skeletal, dental, and soft tissue measures, treatment duration, and patient-reported outcomes across the different treatment groups.

Conclusion

The management of distal occlusion, a prevalent malocclusion characterized by a retruded mandible relative to the maxilla, represents a significant challenge in orthodontics. This review explored various treatment modalities and highlighted the key principles and considerations for optimizing outcomes in these patients. The diverse nature of distal occlusion necessitates individualized treatment planning, with a thorough diagnosis and assessment being paramount for success. From growth modification in children and adolescents to complex surgical-orthodontic approaches

in adults, the goal remains the achievement of a stable, functional, and esthetically pleasing occlusion.

The effectiveness of early interceptive treatment during the growth period is crucial in influencing skeletal discrepancies. Functional appliances, such as the Herbst or Twin Block, have demonstrated the ability to stimulate mandibular growth and reduce the severity of distal occlusion in growing patients. While the success of these appliances is not guaranteed, their early application appears advantageous in milder cases and can simplify treatment in later stages. Careful monitoring of growth and skeletal changes is essential to ensure optimal results and minimize potential side effects.

For individuals with severe skeletal discrepancies, particularly adults with limited or no growth potential, the combined surgical-orthodontic approach offers the most effective method for addressing the underlying skeletal imbalance. Orthognathic surgery, typically involving mandibular advancement, not only corrects the skeletal malocclusion but also improves facial aesthetics. However, this approach requires meticulous planning, coordination between the orthodontist and surgeon, and consideration of the risks and benefits. In such instances, patient compliance with the treatment protocols and clear communication are vital for achieving satisfactory long-term results.

The role of fixed orthodontic appliances, such as braces, remains essential in the refinement and detailing of occlusal relationships. The precision and control afforded by modern bracket systems have greatly contributed to enhancing the efficiency and accuracy of treatment outcomes. Aligners are also gaining popularity for their aesthetic appeal and convenience, and can be effective in managing less severe distal occlusions and achieving predictable outcomes. Careful consideration of the specific malocclusion and patient compliance is essential to determine the appropriate appliance selection.

The importance of long-term stability cannot be overstated. Retention protocols, including the use of fixed and removable retainers, are crucial for maintaining the achieved occlusion and preventing relapse. Patients must be educated

about the importance of compliance with retention protocols to preserve the treatment outcomes for the long-term.

In summary, the optimization of orthodontic treatment for distal occlusion requires a nuanced and individualized approach. Early intervention with growth modification, precise fixed or removable mechanics, consideration of surgical options for severe cases, and diligent retention protocols are all necessary to achieve long-term success. Future research should focus on further refining these techniques, exploring novel materials and technologies, and developing more precise diagnostic tools. An emphasis on patient-centered care, detailed treatment planning, and continuous monitoring will ultimately contribute to achieving more predictable and stable outcomes for patients with distal occlusion. Ultimately, orthodontics should focus on providing stable, functional and esthetic outcomes for every individual.

REFERENCES

- 1.Khalimova K. M. et al. Clinical neurological and diagnostic aspects of headaches with congenital precerebral angiodyplasia //Toshkent tibbiyot akademiyasi axborotnomasi. – 2019. – №. 2. – C. 132-134.
- 2.Salokhiddinov M., Rahmatullaeva G. The role of pathological deformations of the main arteries of the brain In the development< bold> of stroke</bold> //CEREBROVASCULAR DISEASES. – ALLSCHWILERSTRASSE 10, CH-4009 BASEL, SWITZERLAND : KARGER, 2018. – T. 45. – C. 473-473.
- 3.Rahmatullaeva G. Molecular Genetic Features Of Polymorphism Of The Gene Col4a1 Rs605143 G/A In The Development Of Pathological Deformations Of Cerebral Vessels In Patients With Migraine (P2. 241) //Neurology. – 2018. – T. 90. – №. 15_supplement. – C. P2. 241.
- 4.Khalimova K., Rakhmatullaeva G., Rashidova N. Role of the insolventy connective tissue in the development of pathological deformation of cerebral vessels in patients with primary headaches //Journal of the Neurological Sciences. – 2013. – T. 333. – C. e512-e513.