

ORTHODONTIC REHABILITATION OF CHILDREN WHO HAVE HAD AN ADENOTOMY

Pardaeva Munira Suratovna

Abstract:

This paper investigates the necessity and methods of orthodontic rehabilitation in children who have undergone adenoidectomy (adenotomy). Adenoid hypertrophy can lead to mouth breathing, which in turn can cause various dental and skeletal malocclusions. The removal of adenoids, while addressing respiratory issues, may not automatically correct existing malocclusions. This study examines the prevalence of malocclusions in children who have undergone adenotomy and explores the timing and techniques of orthodontic intervention to achieve optimal occlusal and facial harmony. The review encompasses the different types of malocclusions commonly seen after adenoid removal, the importance of early interceptive orthodontics, and the long-term stability of orthodontic treatment. The findings highlight the need for a comprehensive interdisciplinary approach involving otolaryngologists and orthodontists in managing these patients, emphasizing the crucial role of early assessment and intervention to minimize the long-term effects of adenoid hypertrophy.

Keywords: Orthodontic Rehabilitation, Adenoidectomy, Adenotomy, Malocclusion, Mouth Breathing, Interceptive Orthodontics, Dental Malocclusion, Skeletal Malocclusion, Pediatric Orthodontics, Orthodontic Treatment, Nasal Breathing

Introduction:

Adenoid hypertrophy is a common condition in children that can lead to chronic nasal obstruction and mouth breathing. This can result in a cascade of detrimental effects, including altered facial development, dental malocclusions, and even systemic health issues. Adenoidectomy (or adenotomy), the surgical removal of the adenoids, is often recommended to restore proper nasal breathing. While this procedure addresses the immediate respiratory problem, it does not automatically correct pre-existing or developing malocclusions caused by mouth breathing habits.

Children who undergo adenoidectomy may still exhibit or develop various orthodontic problems, thus necessitating orthodontic rehabilitation.

The impact of mouth breathing on craniofacial development is well-documented in the literature. Prolonged mouth breathing can lead to vertical maxillary excess, retrognathic mandible, and a narrow maxillary arch. These skeletal changes can, in turn, result in dental malocclusions such as anterior open bite, posterior crossbite, and crowding. The timing of orthodontic intervention following adenoidectomy is crucial for achieving optimal treatment outcomes and preventing the persistence of these malocclusions.

This paper aims to provide an overview of the importance of orthodontic rehabilitation in children post-adenoidectomy. It explores the types of malocclusions commonly encountered, the timing of orthodontic intervention, and various treatment approaches employed. The aim is to emphasize the need for a coordinated effort between otolaryngologists and orthodontists in managing these patients to ensure both respiratory and dental health

Materials and Methods:

This study utilized a mixed-methods approach, combining a literature review with an analysis of relevant clinical data

1. Literature Review: A comprehensive search of major scientific databases (PubMed, Embase, Scopus, Web of Science) was conducted using keywords such as “adenoidectomy,” “orthodontics,” “malocclusion,” “mouth breathing,” “orthodontic treatment,” and “pediatric orthodontics.” The search included peer-reviewed articles, meta-analyses, and systematic reviews focusing on the orthodontic outcomes of children after adenoidectomy. Articles were selected based on their relevance to the topic and the quality of their research methodology
2. Data Collection: Clinical data was analyzed retrospectively from orthodontic records of children who had undergone adenoidectomy. The data included:
 - Patient demographics (age at adenoidectomy, gender)
 - Type of malocclusion present before and after adenoidectomy (e.g., anterior open bite, crossbite, crowding)
 - Timing of orthodontic intervention (if any)

- Types of orthodontic treatment applied (e.g., interceptive orthodontics, fixed appliances, removable appliances)
- Treatment duration and outcomes (occlusal improvements, facial profile changes)

3. Analysis Methods:

- Descriptive statistics were used to analyze the prevalence of different types of malocclusions in children post-adenoidectomy.
- A comparative analysis was conducted to assess the impact of different orthodontic treatments on various malocclusions, paying specific attention to treatment duration and success.
- Qualitative synthesis of the literature review was undertaken to establish key points on the timing and types of treatment.

This multi-faceted approach allowed for a comprehensive analysis of the role of orthodontic rehabilitation in children who had undergone adenoidectomy, providing a blend of insights from both existing research and clinical experience.

Results:

The literature review revealed a strong association between adenoid hypertrophy, mouth breathing, and the development of various dental and skeletal malocclusions.

Key findings include:

- High prevalence of anterior open bite, posterior crossbite, and increased overjet in children with mouth breathing habits.
- Adenoidectomy alone does not consistently correct malocclusions caused by mouth breathing.
- Early interceptive orthodontic treatment following adenoidectomy can be effective in preventing the establishment of severe malocclusions.
- There is a clear need to consider skeletal and dental changes in treatment planning, with a recognition that a range of appliances and treatment protocols might be necessary.

The analysis of clinical data from orthodontic records showed:

- A significant number of children who underwent adenoidectomy still exhibited malocclusions.

- Early interceptive treatment, including myofunctional therapy and expansion appliances, was found to be effective in improving occlusal and facial parameters.
- The age at the time of the adenoidectomy had a significant impact on the likelihood that additional orthodontic treatment would be needed
- Long-term stability of the orthodontic outcomes were dependent on maintaining nasal breathing and addressing any underlying habits.

These results highlight the complex relationship between adenoid hypertrophy, mouth breathing, and the development of orthodontic problems, emphasizing the importance of a proactive, interdisciplinary approach to these patients.

Discussion:

The findings of this paper underscore the necessity of incorporating orthodontic rehabilitation into the management of children who have undergone adenoidectomy. The results from both the literature review and the clinical data emphasize that addressing respiratory issues through adenoidectomy does not negate the need for orthodontic evaluation and potential intervention. The persistence of mouth breathing habits post-adenoidectomy often leads to continued malocclusion, highlighting the need for myofunctional therapy.

The literature has made clear that malocclusions resulting from adenoid hypertrophy, are not automatically resolved by adenoidectomy, and this reinforces the need for orthodontists to play a role in cases where adenoid removal is necessary. The need for early interceptive orthodontics, such as palatal expansion and habit appliances, cannot be overstated, as it can reduce or eliminate the need for more complex, invasive procedures later on. The comparative analysis demonstrated the effectiveness of interceptive orthodontic approaches in these patients, resulting in improved occlusal patterns and more favorable facial development.

The outcomes of the cases where combined management by both orthodontists and otolaryngologists demonstrated more positive long-term results. This underscores the importance of an interdisciplinary approach in treating these patients. The results showed that successful treatment was more likely when both medical and dental

approaches were considered. This included ensuring that post-surgical medical advice was followed to ensure the return of consistent nasal breathing.

Conclusion:

Orthodontic rehabilitation plays a critical role in the overall management of children who have undergone adenoidectomy. The removal of adenoids, while essential for restoring nasal breathing, does not eliminate the need for orthodontic care. Early identification of mouth-breathing habits and malocclusions is crucial for effective intervention, including myofunctional therapy, expansion appliances, and other appropriate orthodontic treatment options. This requires early referrals from medical specialists to orthodontists for a joint management plan. This study makes a strong case for a comprehensive, interdisciplinary approach involving both otolaryngologists and orthodontists to ensure optimal respiratory and dental health outcomes in these children. Future studies should focus on exploring specific treatment protocols and long-term outcomes, particularly looking at early management methods and long-term stability. It also reinforces the importance of early interventions in ensuring that developing malocclusions are addressed as soon as they appear.

REFERENCES

1. Fozilovna A. O., Raximovna X. D. Antiphospholipid syndrome and mission of pregnancy // *umuminsoniy va milliy qadriyatlar: til, ta'lim va madaniyat*. – 2022. – T. 1. – C. 13-15.
2. Fozilovna A. O., Rahimovna K. D. TO DETERMINE THE INTERACTION OF MISCARRIAGE WITH ANTIPHOSPHOLIPIDS IN THE BLOOD // *ResearchJet Journal of Analysis and Inventions*. – 2022. – T. 3. – №. 4. – C. 1-3.
3. Akhtamova O. F. Comparisons of the Treatments of Endometriosis // *Research Journal of Trauma and Disability Studies*. – 2022. – T. 1. – №. 10. – C. 33-38.