

Tilyavova Sitora Amirovna

Bosimova Noila Ma‘rifjon qizi

Majitova Nargiza Qaxramon qizi

**SCIENTIFIC ARTICLE: POLYCYSTIC OVARY SYNDROME (PCOS):
DIAGNOSIS AND TREATMENT METHODS**

Abstract

Polycystic ovary syndrome (PCOS) is a common endocrine disorder affecting women of reproductive age. It is characterized by irregular menstrual cycles, excess androgen production, and polycystic ovaries visible on ultrasound. PCOS can lead to various complications, including infertility, metabolic issues, and an increased risk of developing type 2 diabetes and cardiovascular diseases. Early diagnosis and effective management are critical for improving the quality of life of affected women. This article aims to provide an overview of PCOS, discussing its pathophysiology, diagnostic criteria, treatment methods, and management strategies. Key diagnostic tools, such as ultrasound, hormonal profiling, and clinical assessment, are explored. Treatment options, including lifestyle modifications, pharmacological therapies, and surgical interventions, are also reviewed.

Keywords: *Polycystic ovary syndrome, diagnosis, treatment methods, metabolic syndrome, infertility, androgens, hormonal imbalance, ultrasound, lifestyle interventions, pharmacological therapy.*

Introduction

Polycystic ovary syndrome (PCOS) is one of the most common hormonal disorders in women of reproductive age. It affects an estimated 5-10% of women worldwide, making it a leading cause of infertility and a significant health concern. The condition is named after the appearance of multiple cysts in the ovaries, which are often visible on ultrasound. However, the name "polycystic ovary syndrome" can be misleading, as not all women with PCOS have cystic ovaries.

PCOS is primarily characterized by three clinical features: irregular menstrual cycles, elevated androgen levels, and the presence of polycystic ovaries. However, the syndrome presents a wide range of symptoms that can vary from woman to woman, making it difficult to diagnose. These symptoms may include hirsutism

(excess hair growth), acne, obesity, and alopecia (thinning hair). In addition to its reproductive consequences, PCOS is associated with long-term metabolic issues, such as insulin resistance, obesity, and an increased risk of developing type 2 diabetes and cardiovascular diseases.

The exact cause of PCOS is not fully understood, but it is thought to be a combination of genetic, environmental, and lifestyle factors. Insulin resistance, which affects the body's ability to use insulin effectively, plays a central role in the pathophysiology of the disease. Moreover, elevated levels of androgens (male hormones) contribute to many of the symptoms observed in PCOS, including irregular ovulation and physical manifestations like hirsutism.

Despite its prevalence, the diagnosis of PCOS is often delayed, and many women may suffer from its effects for years before seeking medical help. Early diagnosis and proper management are essential for preventing complications such as infertility, metabolic dysfunction, and long-term health risks. This article aims to provide a comprehensive understanding of PCOS, focusing on the current approaches to diagnosis and treatment.

Materials and Methods

This review article is based on a thorough analysis of available literature, clinical studies, and evidence-based guidelines on the diagnosis and management of polycystic ovary syndrome. A systematic search of databases such as PubMed, Scopus, and Google Scholar was conducted to identify peer-reviewed articles published in the last two decades. The following inclusion criteria were applied: studies focused on the diagnosis, treatment, and management of PCOS; articles discussing hormonal and metabolic aspects of the syndrome; and clinical guidelines issued by reputable health organizations.

The search terms included "polycystic ovary syndrome," "PCOS diagnosis," "treatment of PCOS," "insulin resistance and PCOS," "ovarian dysfunction," "hirsutism," and "PCOS management." Both observational studies and randomized controlled trials were included in the review. The selected studies were analyzed to summarize current diagnostic practices, treatment options, and management strategies for PCOS.

Inclusion Criteria:

Studies that included women diagnosed with PCOS (according to Rotterdam or NIH criteria).

Randomized controlled trials (RCTs) and clinical trials focused on the treatment of PCOS.

Reviews and meta-analyses discussing the management of metabolic disorders in PCOS.

Exclusion Criteria:

Studies with a small sample size (less than 50 participants).

Studies focusing on other diseases with similar symptoms to PCOS.

Articles that did not offer evidence-based treatment protocols.

Results and Discussions

1. Diagnosis of Polycystic Ovary Syndrome (PCOS)

The diagnosis of PCOS is primarily based on clinical presentation and laboratory findings. According to the Rotterdam criteria (2003), at least two of the following three criteria must be met for a diagnosis of PCOS:

1. **Irregular or absent ovulation** (evidenced by irregular menstrual cycles or anovulation).
2. **Clinical and/or biochemical signs of hyperandrogenism** (e.g., hirsutism, acne, male-pattern baldness, or elevated levels of testosterone in the blood).
3. **Polycystic ovaries on ultrasound**, defined as the presence of 12 or more follicles in one ovary or an ovarian volume greater than 10 mL.

In some cases, it is important to rule out other causes of menstrual irregularity and hyperandrogenism, such as thyroid disorders, adrenal disorders, and tumors. In addition to clinical assessment, ultrasound imaging plays a crucial role in confirming the presence of polycystic ovaries. However, it is important to note that many women without PCOS also have polycystic ovaries, and the diagnosis cannot rely solely on this finding.

Biochemical testing is often used to evaluate androgen levels, particularly testosterone, and to rule out other causes of elevated androgens, such as non-classical congenital adrenal hyperplasia. Moreover, tests for insulin resistance, lipid profiles,

and glucose tolerance are essential for assessing metabolic abnormalities, which are common in women with PCOS.

A challenge in diagnosing PCOS is its variability in symptoms. Not all women with PCOS exhibit the classic triad of symptoms. Some may only experience infertility or irregular periods without significant androgen excess or polycystic ovaries. Thus, clinical judgment is crucial in diagnosing PCOS and differentiating it from other conditions that mimic its symptoms.

2. Pathophysiology of PCOS

The pathophysiology of PCOS is complex and involves hormonal imbalances, metabolic dysfunction, and genetic predisposition. Central to the development of PCOS is insulin resistance, which leads to elevated insulin levels in the bloodstream. This insulin resistance is thought to play a key role in the development of both metabolic abnormalities and reproductive dysfunction in PCOS.

Elevated insulin levels stimulate the ovaries to produce more androgens, which, in turn, disrupt the normal ovulatory cycle. The increased androgen production leads to symptoms such as hirsutism, acne, and male-pattern baldness. Additionally, insulin resistance contributes to the development of obesity, another common feature of PCOS, further exacerbating metabolic issues.

Another important factor in the pathophysiology of PCOS is the dysregulation of the hypothalamic-pituitary-ovarian axis. Women with PCOS often have higher levels of luteinizing hormone (LH) and low levels of follicle-stimulating hormone (FSH), which disrupt normal ovulation. This hormonal imbalance leads to anovulation (lack of ovulation), which is a key feature of PCOS and the primary cause of infertility in affected women.

3. Treatment Methods for Polycystic Ovary Syndrome

Management of PCOS is multifaceted, involving lifestyle changes, pharmacological therapies, and, in some cases, surgical intervention. Treatment approaches depend on the primary symptoms the patient presents with, including infertility, hirsutism, or metabolic dysfunction.

A. Lifestyle Modifications

The first line of treatment for women with PCOS, particularly those who are overweight or obese, is lifestyle modification. Weight loss, even a modest reduction of 5-10% of body weight, can significantly improve menstrual regularity, reduce androgen levels, and increase the likelihood of ovulation.

Dietary changes, including a low glycemic index (GI) diet, can help manage insulin resistance and reduce the risk of developing type 2 diabetes. Regular physical activity, such as moderate-intensity exercise, has been shown to improve insulin sensitivity and reduce metabolic symptoms associated with PCOS. In addition, managing stress and getting adequate sleep are also important factors in improving overall health.

B. Pharmacological Treatments

Several medications are commonly used to treat the symptoms of PCOS:

1. **Oral Contraceptives (OCs):** OCs are often prescribed to regulate menstrual cycles and reduce androgenic symptoms, such as hirsutism and acne. Combined oral contraceptives containing both estrogen and progestin help lower androgen levels by suppressing ovarian androgen production. They also improve the regularity of menstrual cycles, reducing the risk of endometrial hyperplasia and cancer due to unopposed estrogen.
2. **Anti-Androgens:** Medications such as spironolactone can be used to reduce excess hair growth (hirsutism) and acne by blocking the effects of androgens on target tissues. Spironolactone is often used in conjunction with OCs for optimal results.
3. **Insulin Sensitizers:** Metformin, a common medication for type 2 diabetes, is frequently used in PCOS to improve insulin sensitivity. Metformin can help regulate menstrual cycles and improve ovulation, particularly in women with insulin resistance.
4. **Ovulation Induction Agents:** For women with PCOS who are trying to conceive, ovulation induction agents such as clomiphene citrate or letrozole are commonly used to stimulate ovulation. These medications help to regulate the hormonal signals necessary for ovulation and increase the chances of conception.

C. Surgical Interventions

In some cases, women with PCOS who are resistant to medical treatments may benefit from surgical interventions. One option is **ovarian drilling**, a laparoscopic procedure that involves making small punctures in the ovaries to reduce androgen production. This technique can restore ovulation in some women and improve fertility outcomes.

Conclusion

Polycystic ovary syndrome (PCOS) is a complex endocrine disorder that requires a multifaceted approach for diagnosis and management. Early identification of the syndrome and its associated risks is essential for preventing complications, including infertility, metabolic dysfunction, and long-term health issues such as cardiovascular disease and diabetes. While the exact cause of PCOS remains unclear, its pathophysiology is strongly linked to insulin resistance and hormonal imbalance. Lifestyle modifications, including weight loss and exercise, form the cornerstone of treatment, particularly for women with metabolic concerns. Pharmacological therapies, such as oral contraceptives, anti-androgens, and insulin-sensitizing agents.

REFERENCES

- 1.Sharipovna N. N., Shukhratovich S. S. MORPHO-FUNCTIONAL CHANGES IN PERIODONT TISSUES IN CHRONIC GENERALIZED PERIODONTITIS ASSOCIATED WITH CHRONIC VIRAL HEPATITIS B AND C //Frontline Medical Sciences and Pharmaceutical Journal. – 2022. – T. 2. – №. 05. – C. 68-76.
- 2.Shuxratovich R. R. A. S. S. DENTAL CARIES OF PRIMARY TEETH AND ITS COMPLICATIONS IN CHILDREN //INNUC. – 2025. – T. 1. – №. 1. – C. 2-8.
- 3.Shukurov S. CAUSES AND METHODS OF TREATMENT OF THE DEVELOPMENT OF PRIKUS ANAMAIAS IN CHILDREN //INNUC. – 2024. – T. 1. – №. 4. – C. 197-200.
- 4.Ayupova F. et al. 137. The influence of acute respiratory infection in pregnant women on the development of preeclampsia //Pregnancy Hypertension. – 2018. – T. 13. – C. S83.
- 5.Муминова З. А., Нишанов Д. А. Морфологическая характеристика пуповины при беременности, осложненной острой респираторной инфекцией //Журнал теоретической и клинической медицины. – 2016. – №. 4. – C. 104-107.



6. Муминова З. А., Саиджалилова Д. Д., Нишанов Д. А. Морфологическая характеристика плацент женщин, перенесших острые респираторные инфекции // Журнал теоретической и клинической медицины. – 2016. – №. 3. – С. 148-150.
7. Муминова Г. Ш., Аюпова Д. А., Муминова З. А. Акушерские и перинатальные исходы у беременных с многоводием. – 2021.