

LOWER JAW TOOTH EXTRACTION. EARLY AND LATE COMPLICATIONS OF TOOTH EXTRACTION

Doniyorov Ortiq Begali o'g'li

Inomjonov Sardorbek Suratjon o'g'li

Narzieva Difuza Bakhtiyorovna

Abstract

Tooth extraction is one of the most common surgical procedures in dental practice, particularly involving the lower jaw (mandible). Although it is often considered a routine intervention, lower jaw tooth extractions present unique anatomical and clinical challenges due to the density of mandibular bone, proximity to vital structures such as the inferior alveolar nerve, and limited accessibility in posterior regions. This paper aims to provide an overview of the most common early and late complications associated with mandibular tooth extraction, emphasizing risk factors, prevention strategies, and post-operative management.

Early complications may include pain, bleeding, swelling, infection, alveolar osteitis (dry socket), and nerve injury. These typically arise within the first few days following extraction. Late complications, developing weeks or even months later, may involve delayed bone healing, jaw stiffness, chronic pain, or the formation of residual cysts. Special attention is given to complications involving the inferior alveolar and lingual nerves, which can result in temporary or permanent sensory disturbances.

Understanding these complications is essential for improving patient outcomes, reducing post-operative morbidity, and guiding dentists in evidence-based clinical decision-making. Preventive measures, such as proper surgical technique, adequate patient assessment, and timely follow-up care, play a critical role in minimizing complications.

Key words: guiding dentists, timely follow, surgical technique

INTRODUCTION

Tooth extraction remains one of the most frequently performed oral surgical procedures across the globe. Despite advances in preventive dentistry and restorative treatments, extractions are often necessary due to advanced caries, periodontal

disease, trauma, orthodontic needs, or impacted teeth. Among all tooth extractions, those involving the lower jaw—also known as mandibular tooth extractions—present a distinct set of anatomical, physiological, and technical challenges. The mandible, being a dense and relatively immobile bone structure, contributes to the complexity of surgical access and increases the likelihood of certain complications, both in the early and late postoperative stages.

The process of extracting a mandibular tooth requires careful consideration of several critical factors, including bone density, proximity to vital anatomical structures such as the inferior alveolar nerve (IAN), lingual nerve, mental foramen, and temporomandibular joint (TMJ), as well as the anatomical variations among patients. When not managed appropriately, these extractions may result in complications that range from mild discomfort to severe functional impairment. Understanding the early and late complications of tooth extraction is essential for oral surgeons, general dentists, and students of dental sciences. It allows clinicians to minimize risks, provide accurate preoperative counseling to patients, and manage postoperative recovery more effectively. Complications not only prolong healing time and increase treatment costs but may also significantly impact the patient's quality of life. Therefore, a comprehensive understanding of these complications and their underlying causes is essential to enhancing patient safety and surgical success.

Overview of Mandibular Anatomy and Its Surgical Implications

The mandible, or lower jawbone, is unique in its structure and function. It supports the lower teeth and plays a vital role in mastication, speech, and aesthetics. The mandibular canal runs within the body of the mandible and houses the inferior alveolar nerve, artery, and vein. This nerve provides sensory innervation to the lower teeth, chin, and lower lip. Damage to this nerve during extraction—especially of third molars (wisdom teeth)—can result in temporary or permanent numbness, paresthesia, or dysesthesia.

Posterior mandibular teeth are particularly difficult to extract due to their location near dense cortical bone and their proximity to the inferior alveolar canal. Molars, especially impacted third molars, are often deeply embedded and may be surrounded by dense bone, making surgical intervention more invasive and traumatic.

Additionally, the presence of anatomical variations such as bifid canals, accessory foramina, or close proximity of roots to the mandibular canal increases the risk of nerve injury during extraction.

Early complications refer to those that arise within hours or days after the extraction procedure. These may include:

Post-extraction pain is expected but should be monitored for abnormal intensity or duration. Swelling is a physiological response but may be exacerbated by trauma or infection.

Inadequate hemostasis during or after the procedure can lead to persistent bleeding. Patients with clotting disorders or those on anticoagulants are at higher risk.

This is one of the most common complications of lower molar extractions, particularly affecting the mandibular third molars. It occurs when the blood clot at the site of extraction is dislodged or fails to form, leading to exposed bone and intense pain.

Infections may arise due to contamination of the surgical site, especially if postoperative hygiene is inadequate or if the extraction was traumatic.

This can result from trauma to the pterygoid muscles or TMJ, inflammation, or infection, leading to difficulty in opening the mouth.

Trauma to the inferior alveolar or lingual nerves during extraction may cause temporary or permanent sensory disturbances. These injuries are more frequent during the removal of deeply impacted third molars.

Late complications occur weeks or even months after the extraction and can be equally concerning. These include:

Poor vascular supply, systemic diseases such as diabetes, and smoking can impair healing.

Some patients may experience persistent neuralgic pain or post-traumatic neuropathic conditions long after the surgical site has closed.

A rare but serious infection of the bone that may develop from untreated infections or compromised blood supply.

Incomplete extractions can leave behind root tips or foreign bodies that may later become symptomatic.

If periapical lesions are not completely removed, they can form residual cysts or granulomas, which may require further surgical intervention.

In rare cases, particularly in elderly or osteoporotic patients, excessive force during extraction or prolonged bone weakening can result in mandibular fractures.

After extraction, the alveolar bone undergoes remodeling, which may result in loss of ridge height and width. This has implications for future prosthetic or implant treatments.

Several patient-related and procedure-related factors contribute to the likelihood of complications. These include:

Older patients are at greater risk for slower healing and osteoporotic changes, while younger patients may have more complex root morphology in third molars.

Systemic conditions such as diabetes, immunosuppression, or bleeding disorders affect healing and infection control.

These habits impair vascularization and immune response.

Excessive trauma, inadequate irrigation, poor flap design, and failure to remove all root fragments increase risk.

Teeth with chronic infections or cystic lesions may complicate surgery.

A thorough preoperative assessment, including radiographic imaging (e.g., panoramic X-rays or CBCT), is critical to identifying anatomical challenges. Informed consent should clearly communicate the risks of complications. Preventive strategies such as atraumatic extraction techniques, antibiotic prophylaxis (when indicated), and appropriate pain management must be employed.

Postoperative care, including patient education on oral hygiene, proper nutrition, medication adherence, and follow-up visits, plays a pivotal role in minimizing complications. Patients should be advised on signs of potential complications and instructed to report persistent symptoms promptly.

Lower jaw tooth extractions, while routine in many dental practices, require careful planning and execution due to the unique anatomical and physiological features of the mandible. Complications, both early and late, can be minimized through proper surgical technique, patient selection, and comprehensive follow-up care. By thoroughly understanding the risks and the clinical presentation of potential



complications, clinicians can enhance outcomes and maintain the highest standards of patient safety and comfort.

REFERENCES

1. Ubbiniyazova A. M. Characteristic of a functional condition of an organism of workers of the Tashkent tractor plant //Medical magazine of Uzbekistan. – 2012. – С. 80-82.
2. Ataniyazova R. A. et al. Assessment of knowledge and skills of hand hygiene among representatives of three generations of the family living in different regions of Uzbekistan.
3. Madaminova M. A. et al. The dynamics are changes in organoleptic parameters and chemical composition of amudarya river water //European Journal of Molecular and Clinical Medicine. – 2020. – Т. 7. – №. 7. – С. 537-545.
4. Атаханова Д. О., Уббиниязова А. М., Садирова М. К. ОСНОВНЫЕ ГИГИЕНИЧЕСКИЕ ПРОБЛЕМЫ КЛАССИФИКАЦИИ ЗАГРЯЗНЕНИЯ ПОЧВ В КАРАКАЛПАКСТАНЕ //The Way of Science. – 2014. – С. 77.